



Private Forests Tasmania

Farm Forestry Series - Eucalypts



This series on Eucalypt Farm Forestry has five parts. Each part can be read individually or as part of the series.

1. Overview
2. Establishment
3. Clearwood Production
4. Pulpwood Production
5. Cost and Labour Estimates

OVERVIEW

Tasmania's premium native forest eucalypts for timber production include *Eucalyptus regnans* (Mountain ash), *E.delegatensis* (White topped stringybark) and *E.obliqua* (Brown stringybark). These three species are members of the ash group and marketed collectively as Tasmanian Oak. Their prime use is appearance grade applications such as furniture, flooring and veneers, with pulpwood production a by product from low quality logs. The species has performed poorly when grown in plantations. *E.globulus* (southern blue gum) is another common Tasmanian eucalypt, generally from lower elevation sites in south east Tasmania. It is a dense, strong and relatively durable timber that has been used extensively for heavy construction and flooring and is an excellent pulpwood species. It is a very successful plantation species, having become the most widely planted eucalypt in the world. Growth and form are excellent on suitable sites, although frost resistance is only moderate. *E.nitens* (shining gum) is the most common plantation eucalypt species in Tasmania. In its natural environment it is primarily a sub-alpine species from high elevation sites in eastern Victoria and NSW, with excellent frost resistance, growth and form. It is a member of the southern blue gum group that includes *E.globulus*, although timber properties are more similar to the ash eucalypts. Uses include construction, flooring, joinery and pulpwood.

Mature Forest

Mature forests are preferable for solid timber production and veneers as the trees have attained large dimensions with a large proportion of mature, stable timber.

Regrowth Forest

Regrowth forests, particularly those less than ~80 years of age, are often inferior to mature forests for solid timber production. Trees contain a greater proportion of young, juvenile timber that has lower density, strength and durability than do mature trees. Regrowth trees may have relatively high growth stresses, potentially resulting in some difficulties with sawing and drying. Smaller diameters lead to a lower output of higher quality timber due to defects such as knots, while regrowth forests as a whole usually have a lower proportion of sawlogs and veneer logs. They do however, produce timber with a higher quality pulp yield than do mature forests.

EUCALYPT PLANTATIONS

Pulpwood Production

E.nitens and *E.globulus* are primarily grown for pulpwood production on short rotations of 12-20 years, depending upon site quality. *E.globulus* produces a higher quality pulp than *E.nitens*, although *E.nitens* is more widely grown in Tasmania due to greater frost tolerance and productivity, particularly on colder, higher elevation sites. These plantation eucalypts produce a higher grade of pulpwood than mature native forests.

Sawlog Production (Clearwood)

The growing of plantation *E.globulus* and *E.nitens* for sawlog and veneer log is still being developed. *P.radiata*



Private Forests Tasmania

Farm Forestry Series - Eucalypts



is a proven plantation species for appearance grade products with well developed markets, to date this is not the case with *E.globulus* and *E.nitens*. While plantation grown clearwood logs have been sawn and veneers sliced, such processing has been largely experimental, with mixed results. Silvicultural regimes for producing clearwood with acceptable wood properties for appearance grade products are being refined. Evidence suggests that high quality logs may be produced on rotations of 20-25 years or more, depending upon site productivity. It is becoming apparent that to grow high quality clearwood, attention to pruning, low stocking rates from a young age and large diameter trees are required to minimise growth stress and the formation of tension wood.

Density & Durability

E.globulus has a dry density of ~900 kg/m³ and a durability rating of class 3.

E.nitens has a dry density of ~700 kg/m³ and a durability rating of class 4.

¹The above ratings are for the heartwood of mature trees.

Durability Ratings (general description for untreated timber)

Class 1 - Very durable: life expectancy of 25 years or more when used in the ground, excellent for above ground use in exposed situations.

Class 2 - Durable: expected life of 15-25 years in the ground, good for above ground use in exposed situations.

Class 3 - Moderately durable: generally unsuitable for in ground use, potentially suitable for some exposed situations.

Class 4 - Non-durable: not suitable for in ground use or exposed situations.

Some Other Properties of Eucalypts

Young plantation grown trees can be expected to have a significantly lower density at a very young age. Density, which increases with tree age, is related to timber strength and hardness and the durability of young timber may be less than that of mature timber. Thus, the timber properties of young trees differ from those of mature trees. Heartwood is usually difficult to treat with preservatives. The sapwood of all species is not durable and requires preservative treatment if utilised outdoors. The dry sapwood of many species (including *E.globulus* and *E.nitens*) is also susceptible to insect attack (lyctid borers) and may require chemical treatment. This can be important if trees with a high proportion of sapwood are processed for purposes such as internal framing timbers or flooring. Significant attack by insects can destroy the untreated sapwood of susceptible species.

END USES (Plantation *E.globulus* & *E.nitens*)

- **Pulpwood** - Bell Bay, Hampshire (Burnie) and Triabunna processing plants. Gunns Ltd exports from all 3 locations while Forest Enterprises Australia Ltd exports from Bell Bay.
- **Posts & Poles** - Preservative treatment is required. Limited market due to the size of the plantation resource compared to the local market for posts and poles. Eucalypts offer a strength advantage over *Pinus radiata* for posts with similar diameter and branching characteristics.
- **Firewood** - Very limited market potential. Young plantation trees have significantly less density and heat value than native forest eucalypts traditionally sourced for the firewood market.
- **Small Diameter Sawlogs** - Forest Enterprises Australia operates a Hew Saw at Bell Bay that utilises small diameter, unpruned plantation grown eucalypts. Logs are sourced with a small end diameter of 16cm to a large end diameter of 34cm.
- **Clearwood Sawlogs & Veneer Logs** - With advances in silviculture, sawing techniques and drying schedules for plantation grown sawlogs, it is anticipated that pruned *E.globulus* and *E.nitens* will become



Private Forests Tasmania

Farm Forestry Series - Eucalypts



an established source of appearance grade clearwood. Tree breeding programs in the future may target improved characteristics such as improved density and branching habits, as has been undertaken for *Pinus radiata*.

SITE & SPECIES SELECTION

The following table is a guide to the anticipated site productivity for *E.globulus* and *E.nitens* plantations in Tasmania². The most limiting factor on any site determines the likely productivity class. The Farm Forestry Toolbox can also be used to estimate productivity. The most accurate estimate of productivity can be gained by measuring growth of existing adjacent or nearby plantations with similar land attributes.

LAND ATTRIBUTES	Class 1A MAI >30	Class 1B MAI 20-30	Class 2 MAI 15-20	Class 3 MAI 10-15	Class 4 MAI <10
Productivity	Very High	High	Medium	Low	Very Low
Elevation (<i>E.globulus</i>)	<300 ^a	<300 ^a	300-400 ^b	400-600	>600
Elevation (<i>E.nitens</i>)	<600	<600	600-850 ^c	600-850	>850
Mean Annual Rainfall (MAR) (mm/year)	>1100	>1100	850-1100	700-850	<700
Drainage	Rapid	Moderate	Imperfect	Poor	Very Poor
Soil Profile					
Effective Rooting Depth (ERD)	>80cm <10%	>80cm 10-30%	45-80cm 30-50%	20-45cm 50-90%	<20cm >90%
Stoniness					
Nutrient Status					
Phosphorus - P (ppm)	>250	100-250	100-250	<100	<100
Nitrogen - N (%)	>0.2	0.1-0.2	0.1-0.2	<0.1	<0.1

- a) Limit is lower in colder areas.
- b) Milder coastal areas
- c) Sites with good air drainage, minimal frost severity and snow damage with limited exposure.

Elevation can be used as a guide to the average temperature and length of the growing season (both decrease with increasing elevation) and the incidence and severity of frosts (increased with increasing elevation).

Elevation and frost (frequency and severity) is the major consideration when deciding upon either E.nitens or E.globulus as the preferred plantation species.

Mean Annual Rainfall (MAR) is a measure of moisture availability. It can be affected by other factors such as soil type and depth (water holding capacity), aspect and slope.

Drainage is an indicator to the extent of waterlogging, as it is detrimental to the growth of trees. Ripping and mounding may reduce the extent of waterlogging during the establishment phase.

Soil profile is determined by the Effective Rooting Depth (ERD) and the stone content as a percentage of the soil volume within the ERD zone. ERD is the depth to which a physical barrier impedes deeper root



Private Forests Tasmania

Farm Forestry Series - Eucalypts



development, such as bedrock, solid clay subsoils, waterlogged zones or stone horizons containing little or no fine earth. Deep ripping can shatter compacted subsoils or cemented pans to increase the ERD in some conditions.

Nutrient status can be determined by soil analysis of the top 10cm of mineral soil, although this may not always be a true indication, depending upon soil type. The most widespread limiting nutrients in Tasmania are phosphorus (P) and nitrogen (N), although other nutrients may be limiting in some soil types. Fertilisation can offer a significant increase in growth on many soil types if it is the most limiting factor to site productivity.

Position in the landscape may impact upon the previous land attributes.

- Sites with poor cold air drainage, resulting in lower temperatures and frost hollows, can occur even at relatively low elevations close to the coast.
- Northern aspects will have higher temperatures than southern aspects.
- Southern and westerly aspects usually experience greater wind exposure than northern and eastern aspects.
- The soil profile, nutrient status and moisture availability is often greater at lower positions within the landscape.

PESTS & DISEASES

A number of pests and diseases are known to impact upon plantations of *E.globulus* and *E.nitens*. Most are of minor significance, although some can cause considerable loss of productivity if monitoring and control measures are not undertaken.

Chrysomelid beetles, particularly the Tasmanian eucalypt leaf beetle, can be major insect pests of *E.globulus* and *E.nitens*. Severe defoliation can occur during periods of significant outbreaks. Control can be undertaken with insecticides.

Caterpillars, such as the Autumn gum moth and Gum leaf skeletoniser, can result in significant defoliation. Control can be undertaken with insecticides.

Sap sucking insects, such as coreid bugs can damage leading shoots. Again, control can often be undertaken with insecticides if the level of damage is significant.

Fungal infections can impact upon leaves, stems and roots. The incidence and severity of most diseases are of minor importance.

Browsing animals, such as wallabies, pademelons, rabbits and hares can cause significant damage to seedlings following planting and until the trees attain 1.0-1.5m in height. Brush-tail possums can cause significant browsing damage, including physical damage such as breaking leaders when climbing. Deer may cause problems in areas where populations are known to occur.

Note: The Farm Forestry Toolbox is an excellent source of information and quality pictures concerning pests and diseases of eucalypts in Tasmania. It details the symptoms, significance and known management prescriptions.

ALTERNATIVE EUCALYPT SPECIES

E.globulus and *E.nitens* are not the only eucalypt species currently being grown for clearwood production, although they are the dominant plantation species for pulpwood production in Australia. Other species currently being grown in southern Australia and New Zealand for solid timber production (clearwood) include:



Private Forests Tasmania

Farm Forestry Series - Eucalypts



E.cladocalyx (Sugar gum)

Tolerant of low rainfall and poor soils. Low frost tolerance, with slow to moderate growth rates. Currently milled from older shelterbelts in western Victoria for appearance grade timber (veneers and solid timber) and firewood. Class 1 durability, low growth stress, range of indoor and outdoor uses. Light brown coloured timber.

E.maculata (Spotted gum)

Tolerant of low to moderate rainfall and poor soils. Low frost tolerance, slow to moderate growth rates with good form. Class 2/3 durability, very strong and hard timber, low growth stress, range of indoor and outdoor uses. Light brown coloured timber.

E.tricarpa & *E.sideroxylon* (Red iron bark)

Very tolerant of low rainfall and poor soils, slow growth rates, moderate to high frost resistance. Class 1 durability, very strong and hard timber, low growth stress, wide range of indoor and outdoor uses. Dark red coloured timber. Sapwood rarely attacked by lyctid borers.

E.saligna (Sydney blue gum)

Moderate to high quality soils and rainfall required, moderate to fast growth rates, low frost resistance. Class 3 durability, strong timber, high growth stress, wide range of indoor uses. Red timber that is favoured in New Zealand (trees with interlocked grain sourced for milling to overcome growth stress), although pests and diseases have impacted upon growth and survival in New Zealand.

E.botryoides (Southern mahogany)

Tolerant of moderate quality soils, moderate rainfall and sites with poor drainage. Tolerance of exposure and saline winds, with moderate growth rates, frost resistance and form. Class 2/3 durability, hard, strong timber, can have high growth stress, range of indoor and outdoor uses. Red timber that is favoured in New Zealand, although pests and diseases have impacted upon growth and survival in New Zealand. Sapwood rarely attacked by lyctid borers.

E.muelleriana (Yellow stringybark)

Tolerant of moderate quality soils and rainfall, moderate growth rate and form, low to moderate frost resistance. Class 2/3 durability, hard and strong timber, low growth stress, wide range of indoor and outdoor uses. Increasing favour among New Zealand farm foresters and a favourite of processors due to low growth stress. Small areas planted in NW Tasmania over the last few years with reasonable growth to date on suitable sites. Light brown coloured timber. Excellent ability to recover from fire. Sapwood not susceptible to lyctid borers.

E.globoidea (White stringybark)

Species with very similar tree and timber characteristics to *E.muelleriana*. Limited plantings in Australia, although finding increasing favour in New Zealand.

E.pilularis (Blackbutt)

Species with very similar timber characteristics to *E.muelleriana*. Grown extensively in New Zealand, with good form, although higher growth stress. Also grown in NSW and southern QLD.

NB Many of these alternative species have been planted in Tasmania on a trial basis by Private Forests Tasmania, landowners and other organisations. However, while it is to be encouraged, the planting of these or other alternative species in Tasmania should be on a small scale and considered experimental, as their performance is relatively unknown at this stage. It is possible that some alternative species may fill a niche role, such as the production of naturally durable timber that can be used for exterior use without chemical treatment. It is not expected that these alternative species could be planted on a large scale, but rather as smaller plantings on suitable sites. Thus, they may offer landowners an



Private Forests Tasmania

Farm Forestry Series - Eucalypts



opportunity to enter a high-value niche market that is not dominated by larger industrial scale growers. Landowners should seek further advice regarding species and site selection if considering alternative species.

Hybrid Eucalypts

E.camaldulensis (River red gum) and *E.globulus*

E.camaldulensis is widespread on mainland Australia and is often found in areas of low rainfall and naturally high salinity levels. It has been crossed with *E.globulus* to form a hybrid for planting in areas subject to particularly dry conditions and areas affected by salinity.

E.grandis (Flooded gum) and *E.globulus*

E.grandis is predominantly found in coastal regions of northern NSW and southern QLD. It has been crossed with *E.globulus* to form a hybrid for planting in areas subject to wet soil conditions (waterlogging).

These hybrid eucalypts, although currently expensive at about \$1.80 per seedling, offer the opportunity to plant productive eucalypts on sites that are prohibitive for the growth of *E.globulus* or *E.nitens*. Approximately 40,000 of these hybrids were planted in Tasmania in 2003. Results of these and earlier plantings indicate excellent survival and growth on harsh sites where other eucalypts have failed. Woodlea Nursery of Scottsdale has a license to grow these seedlings in Tasmania.

FURTHER READING

1. Bootle, K.R.(1983) Wood in Australia: Types, properties and uses. McGraw-Hill Book Company, Sydney.
2. Laffan, M.D.(1997) *Site Selection for hardwood and softwood plantations in Tasmania. A methodology for assessing site productivity and suitability for plantations using land resource information.* Soils Technical Report No.3, second edition, Forestry Tasmania and Forest Practices Board, Tasmania.
3. Farm Forestry Toolbox - Free copies are available from Private Forests Tasmania. The Toolbox is a collection of computer programs in a user friendly format on a CD and can assist with the management and planning for both native forest and plantation eucalypts.

For further information contact	
Private Forests Tasmania	
Burnie	6434 6319
Hobart	6233 7445
Launceston	6336 5300
Email: admin@privateforests.tas.gov.au	
© Private Forests Tasmania	
Date	August 2004

This information has been prepared by Private Forests Tasmania. Every reasonable endeavour has been used to ensure that the material was accurate at the time of publication. However, Private Forests Tasmania takes no responsibility for the accuracy, completeness or relevance of such information or for matters arising from changed circumstances or information or material which may have become available subsequently. This information is introductory in nature and should not be treated as a substitute for specific advice or relied on as a basis for business decisions. Before undertaking any significant forestry project it is recommended that you seek personal professional advice directly from Private Forests Tasmania on the particular matter.