



Private Forests Tasmania

Farm Forestry Series - Eucalypts



This series on Eucalypt Farm Forestry has five parts. Each part can be read individually or as part of the series.

1. Overview
2. Establishment
3. Clearwood Production
4. Pulpwood Production
5. Cost and Labour Estimates

PLANTING STOCK/SEED SOURCE

The Southern Tree Breeding Association (STBA) of Australia has developed breeding programs to target above average characteristics of growth and density. Some larger industrial plantation companies have also developed breeding programs for improved growth rates. Seedlings of *E.globulus* and *E.nitens* grown from improved seed sources are available from some Tasmanian nurseries, such as Woodlea Nursery, Scottsdale, ph 6352 7262, Lanoma Forest Nursery, Westerway, ph 6288 1124 or 0418881124 and Forestry Tasmania, Perth, ph 6391 6300.

Table 1 - Seed level and genetic gain for improved seed for *E.globulus* and *E.nitens*¹.

<i>E.globulus</i>		<i>E.nitens</i>	
Level	Growth	Level	Growth
3	10-15	3	10-15
4	15-20	4	15-20
5	20-25	5	20-25
6	25-30		
7	30-35		

NB The above figures for growth are the predicted genetic gain over unimproved native forest seed. The higher the predicted gain the greater the growth rate.

Cost - As the level of seed improvement increases, additional seedling cost is incurred. Indications from nurseries are that level 5 seedlings cost about \$0.05 more than level 3. Levels 6 and 7 (*E.globulus*) are obtained from more intensive breeding programs than lower levels and costs can be expected to rise further.

INITIAL STOCKINGS

Pulpwood Production

As a guide, establishment at spacings of 3.0m x 4.0m (833 stems/ha) to 2.5m x 3.5m (1143 stems/ha) provides an adequate stocking for pulpwood and clearwood production. Industrial scale plantations are typically established at stockings of about 1000 stems/ha.

Stockings at the lower end of the range:

- Require excellent survival to ensure maximum site occupancy and wood production.
- Have slightly lower establishment costs (reduced site preparation costs, seedlings, etc.)
- Have larger piece size at harvest (potentially reduced harvesting costs) for pulpwood regimes.

Stockings at the higher end of the range are recommended for colder, high elevation sites, particularly those subject to exposure. A higher stocking is advisable for clearwood production as it provides a greater number from which to select trees of suitable form, vigour and spacing for clearwood pruning.



Private Forests Tasmania

Farm Forestry Series - Eucalypts



Row spacing should be wide enough (3.5 to 4.0m is recommended) to enable access with equipment such as a tractor for slashing and post-planting weed control. Wider row spacings also reduce cultivation costs when ripping and mounding.

SITE PREPARATION

Site preparation prior to planting is required to ensure successful establishment and involves the following:

- Cultivation (required most sites)
- Weed Control (essential)
- Browsing Control (essential)

CULTIVATION

Soil cultivation is usually required to:

- Increase the effective rooting depth and ease of root penetration through the soil profile.
- Increase moisture availability (particularly at depth), survival and uniform growth.
- Reduce waterlogging during the establishment phase on wet sites by mounding.
- Improve weed control, particularly when residual herbicides are applied to cultivated soil.
- Provide easier and cheaper planting.

Deep Ripping to a depth of 60cm to 1m should be undertaken prior to the autumn break when soils are dry to enable the subsoil to be shattered. Ripping to these depths requires large bulldozers. Deep ripping should not be undertaken when soils are very wet, as the subsoil is sliced rather than shattered. This can lead to soil damage and the concentration of roots along the rip lines, increasing the potential for windthrow.

Note: Ripping is not suitable for all soil types. Deep sands will not benefit from ripping. Soils with dispersible subsoils should not be ripped as additional water infiltration can result in dispersion of the subsoil, leading to potential tunnel and gully erosion.

Mounding concentrates topsoil along the planting lines and mounds should be positioned over the rip line. Mounds should be at least 30cm above the general ground level. Higher mounds are advisable for wet sites to increase drainage and successful establishment. Mounds should not consist of large clods as air pockets can reduce seedling survival. A second pass with cultivation equipment can be useful to reduce the extent of large clods of soil in the mound. Mounding in the autumn followed by spring planting is beneficial to enable winter rains to settle the mounds and reduce air pockets.

Row direction is important to conserve moisture and reduce the potential for soil erosion following cultivation. Mounding on the contour is advisable even on relatively flat sites. On wet sites a slight fall towards drainage lines can help to reduce waterlogging. On dry sites, a slight fall away from the drainage lines towards the ridges can improve moisture retention.

First Rotation Sites

It is standard practice to undertake ripping and mounding for the establishment of eucalypt plantations on ex-pasture sites and ex-forest sites. Cropping in particular can result in soil compaction. Some sites can be planted without cultivation, such as steep slopes with well-structured soils. Poorly structured soils will require cultivation. Spot cultivation may be possible on steeper sites.

Private Forests Tasmania have Savannah mound ploughs available for hire that are designed for farm forestry ex-pasture sites. An 80hp 4WD tractor or larger is suitable for use with this equipment.



Private Forests Tasmania

Farm Forestry Series - Eucalypts



Second Rotation Sites

Second rotation plantation sites are often spot cultivated. Spot cultivation is expensive, with indicative costs of \$850-900/ha. It is possible to plant between the stumps on second rotation sites, and, although planting costs are higher, cultivation expense is foregone, resulting in considerably cheaper establishment. Snig tracks and landings usually require deep ripping to reduce compaction.

Note: The Forest Practices Code 2000 provides guidelines and restrictions upon cultivation practices, depending upon soil type, erosion potential and slopes. Refer to the Forest Practices Code 2000 or seek further advice regarding cultivation for plantation establishment.

WEED CONTROL

Important Note: The use of proprietary names for products is for explanatory purposes only and does not imply any endorsement by Private Forests Tasmania.

(Refer to Private Forests Tasmania's Weed Control Information Sheet for an overview of weed control)

Pre-Planting Weed Control

Successful weed control prior to planting is essential to ensure successful, uniform establishment of eucalypts. The use of residual herbicides in accordance with label directions is particularly important to ensure weed control into the first summer. The presence of weeds during dry summer and autumn conditions can greatly reduce early growth and survival. Failure to control weeds successfully prior to planting can result in major difficulty and expense associated with post-planting weed control.

- Slash or graze the site to reduce the existing weed volume if grasses and broadleaved weeds are the major weed species. Woody weeds such as blackberry and wattles should be treated with herbicides 6-12 months prior to slashing or burning to ensure sufficient uptake and death of the root systems, ensuring minimal germination post-planting.
- Strip or broadscale spray with knockdown herbicides (eg: glyphosate) along planting lines 4-6 weeks prior to cultivation. Although sites can be cultivated without initial knockdown, a finer tilth is often achieved if existing weeds and associated root systems are dead.
- Apply residual herbicide and plant in accordance with the label instructions.

General herbicide options for knockdown of existing weeds

- Glyphosate (eg: Roundup™) - knockdown capability only.
- Brush-Off™ or Garlon™ for the control of hard-to-kill perennial weeds such as blackberry.

General herbicide options for residual control of weeds

- *Eucmix™ Pre Plant* - A water dispersible herbicide for *E.globulus* and *E.nitens* plantations. It has limited knockdown capability. Existing weeds are best controlled prior to application, although Eucmix™ Pre Plant can be mixed with glyphosate. It is designed to control weeds prior to the planting of healthy seedlings taller than 20cm.
- *Lontrel™ Forestry Herbicide* - A water soluble herbicide for use with eucalypts for pre-emergent control of flatweeds, legumes (including wattles), thistles and capeweed. It has no effect on grasses.
- *Simazine* - residual control only.

Note: These residual herbicides have restrictions upon the method and volume of application depending upon soil types, slopes, proximity to watercourses, etc. More effective and longer lasting residual control of weeds is achieved when applied to bare soil following cultivation. Do not cultivate following the application of residual herbicides because soil disturbance will greatly reduce the effectiveness of the herbicide. Read the label carefully and follow any restrictions upon use. Seek expert advice before carrying out on ground work.



Private Forests Tasmania

Farm Forestry Series - Eucalypts



Post-Planting Weed Control

Herbicide options for knockdown and continued residual control of weeds

- *Euclmix GRTM Granular Herbicide* - A granular herbicide that does not need to be mixed with water. For use in *E.globulus*, *E.nitens* and *E.regnans* plantations. It is designed to control weeds less than 10cm in height and diameter. Application is made 12 months after planting. A weed-a-metre can be used for spot application. Sufficient soil moisture is required for uptake by weed species.
- *LontrelTM Forestry Herbicide* - Water soluble herbicide for use in eucalypt plantations for the control of a range of broad-leafed weeds. It has no effect on grasses.

Herbicide options for knockdown of existing weeds

- *VerdictTM* - Selective herbicide for the control of grasses only. Can be sprayed over the top of eucalypts, provided label instructions are followed carefully.
- *Glyphosate* - Shielding against spray contact is essential as eucalypts are extremely sensitive to glyphosate. Can be successful with extreme care, although generally not recommended. Where possible, safer options for weed control are advisable.

Slashing: between the rows reduces competition from weeds and improves access.

- Rows must be wide enough for a tractor and slasher and clear of obstructions.
- Steep side slopes can be hazardous.

Mulching: may also reduce weed competition but is not widely used due to a lack of proven cost competitive techniques.

BROWSING CONTROL

(Refer to Private Forests Tasmania's Farm Forestry Information Sheet - Browsing Damage to Seedlings)

- Eucalypts are palatable to browsing species such as wallabies, possums, rabbits, hares and livestock.
- Maintaining strips of pasture between the planting lines, or undertaking spot weed control, retains an alternative food source. However, browsing should always be anticipated and prevention taken prior to planting.
- Heavy browsing can be expected on ex-forest sites where alternative food sources may be limited.
- Poisoning, shooting and trapping may be used individually or in combination. A permit from the Parks and Wildlife Service is required for these activities.
- Stock-proof fencing is required to exclude cattle and sheep. Stock may browse young trees or cause bark stripping. Cattle can also cause physical damage by pushing trees over as they move through the plantation.
- Although expensive, wildlife-proof fencing to exclude native animals is recommended where browsing pressure is high. Wire netting with an electric outrigger to exclude possums or electric fences with multiple hot-wires can be used.
- Tree guards can be utilised. They are too expensive in most instances and some guard designs can result in poor initial growth due to a lack of ventilation inside the guards.

FERTILISING

(Refer to Private Forests Tasmania's Farm Forestry Information Sheet - Fertilisers)

Fertilising at establishment can substantially increase early growth and survival on most sites.

- As a general rule, 100-150g DAP (Di Ammonium Phosphate) is recommended. Apply 1-2 months post-planting, provided adequate weed control has been achieved.
- Nitrogen (N) is usually the most limiting nutrient for eucalypts, particularly on lower quality sites. Later



Private Forests Tasmania

Farm Forestry Series - Eucalypts



- age fertilising with N can substantially increase growth rates on most sites.
- High quality basalt soils, particularly those with a history of fertiliser input, may not require fertilising at establishment. High N levels can result in poor form due to multiple leaders.
 - Micronutrient (trace element) deficiencies can occur on some sites. DAP containing micronutrients can be applied at planting if the deficient nutrients are known.
 - Additional fertiliser, if required, should be applied at, or before, age 3 to ensure rapid canopy closure and maximum site occupancy.

Identification of fertiliser requirements can be obtained from existing plantations growing on similar soil types in the area where fertiliser requirements may be known. Once the deficient nutrient(s) are identified, fertilising can be undertaken from the ground or by air.

It is recommended that professional advice be obtained, as it may be difficult to determine the exact fertiliser requirements for particular sites.

1. Data sourced from Seed Energy Pty Ltd.

For further information contact
Private Forests Tasmania
Burnie 6434 6319
Hobart 6233 7445
Launceston 6336 5300
Email: admin@privateforests.tas.gov.au
© Private Forests Tasmania
Date August 2004

This information has been prepared by Private Forests Tasmania. Every reasonable endeavour has been used to ensure that the material was accurate at the time of publication. However, Private Forests Tasmania takes no responsibility for the accuracy, completeness or relevance of such information or for matters arising from changed circumstances or information or material which may have become available subsequently. This information is introductory in nature and should not be treated as a substitute for specific advice or relied on as a basis for business decisions. Before undertaking any significant forestry project it is recommended that you seek personal professional advice directly from Private Forests Tasmania on the particular matter.